

**American Society of PeriAnesthesia Nurses**  
**A Position Statement on Waste Anesthesia Gases**  
**Outside of the Operating Rooms**

*Developed by ASPAN and supported by the American Industrial Hygiene Association*

The American Society of PeriAnesthesia Nurses (ASPAN) is the premier organization for perianesthesia nursing excellence. The purpose of ASPAN is to empower and advance the unique specialty of perianesthesia nursing. Perianesthesia registered nurses work in any space before or after patients receive anesthesia, primarily in the postanesthesia care unit (PACU) recovering patients from procedures and surgeries.

**Background**

Established in 1939, the American Industrial Hygiene Association (AIHA) is the association for scientists and professionals committed to preserving and ensuring occupational and environmental health and safety in the workplace and community. More than half of AIHA's 8,500 members are certified industrial hygienists. This organization supports members and those employed in private or public sectors through expertise support, comprehensive education programs, and service as a resource to help maintain the highest workplace and professional standards and for working communities.

The AIHA is composed primarily of industrial hygienists who are trained to anticipate, recognize, evaluate, and recommend controls for environmental and physical hazards that can impact the health and well-being of workers.

ASPAN supports recognition that waste anesthesia gases (WAGs) are an occupational hazard. The National Institute for Occupational Safety and Health (NIOSH) and the Occupational Safety and Health Administration (OSHA) have identified health effects from WAG exposure including nausea, dizziness, headaches, infertility, miscarriages, birth defects, cancer, liver, and kidney diseases.<sup>1-3</sup> The immediate postoperative patient is the source of exposure to WAGs outside of the operating rooms (ORs). Volatile anesthetic gases (e.g., nitrous oxide), and the fluorinated gases (e.g., sevoflurane, desflurane, isoflurane) undergo minimal metabolism in the body during clinical use and are primarily eliminated unchanged via exhalation (estimated equal or greater than 95%).<sup>4,5</sup> Few adverse effects associated with WAGs have been documented when workplace exposure limits are implemented by using measures to reduce occupational exposure.<sup>6</sup> The AIHA supports this ASPAN position statement and stands behind enhancing knowledge and protections to guard nurses from potential occupational hazards faced in the workplace.<sup>7,a</sup>

*<sup>a</sup>The American Industrial Hygiene Association and American Society of PeriAnesthesia Nurses collaborated to produce a white paper on the recognition and control of waste anesthesia gases. To view the document click this link: [Recognition-Evaluation-and-Control-of-Waste-Anesthetic-Gases-in-the-Post-Anesthesia-Care-Unit-White-Paper.pdf](https://www.aspan.org/Recognition-Evaluation-and-Control-of-Waste-Anesthetic-Gases-in-the-Post-Anesthesia-Care-Unit-White-Paper.pdf) (aspan.org)<sup>7</sup>*

## Position

The goals for controlling WAG exposure in hospitals are:

- To increase awareness that all areas of the hospital can have potential occupational exposure to WAGs in concentrations higher than in the operating rooms
- To prevent exposure to WAGs as an occupational hazard for nurses, patients, and their families in all areas of the institution

WAGs should be regulated with *engineering controls*<sup>b</sup> in the PACU including (see Figure 1):

- Application of commercially available source-control scavenging systems to patients recovering in the PACU following surgery, following the administration of inhalation anesthetic agents. Total intravenous anesthesia (TIVA) patients would only require scavenging systems for pathogens
- Increased air-exchanges to provide adequate ventilation similar to operating room air-exchanges, so that WAG levels do not increase as a result of expanding patient volumes
- Provision of fresh air exchanges in the PACU, similar to operating room air recommendations, to reduce and prevent nurse exposure to WAGs

*Administrative controls* for the PACU should include:

- Implementing an industrial hygiene program including routine monitoring of WAGs in the perianesthesia registered nurses' breathing zone and regular maintenance of ventilation and source-control scavenging systems at least twice per year<sup>2,8</sup>
- Developing and administering an education and training program for perianesthesia registered nurses exposed to WAGs that comply with NIOSH Recommended Exposure Limits (RELs), and with OSHA's Hazard Communication Standard<sup>c</sup>
- Defining and implementing best work practices to help reduce perianesthesia registered nurse exposure to WAGs
- Executing a medical surveillance program for all perianesthesia registered nurses exposed to WAGs
- Complying with existing federal, state, and local regulations and guidelines developed to minimize hospital personnel exposure to WAGs
- Encourage and promote the use of patient source-control scavenging systems.

Currently, these controls are not present in the PACUs in which the patients are the source of the WAG as they exhale unmetabolized anesthetic gases into the unregulated environment of the PACU and intensive care areas.

<sup>b</sup>Engineering controls are applied at the source of the hazard and are generally the preferred and most effective means to control and minimize occupational exposures. Administrative controls are the next best means and include implementation of best practices.<sup>2</sup>

<sup>c</sup>OSHA's Hazard Communication Standard requires that healthcare workers are provided information related to workplace chemical hazards.<sup>2</sup>

Relatively recent findings have shown that patients in the PACU can be significant sources of WAG exposure to nurses.<sup>9-12</sup>




Figure 1. Hierarchy of Controls. Reprinted with permission from the National Institute of Occupational Safety & Health (NIOSH).

<https://www.cdc.gov/niosh/topics/hierarchy/default.html>

This position statement serves as the basis for ASPAN’s action on behalf of industrial hygienists, perianesthesia registered nurses, healthcare providers in the perianesthesia environment, and those whom the organizations serve.

## Comments

### 1. Health Outcomes Related to Waste Anesthesia Gases (WAG)

For more than 50 years, the evidence presented in the peer-reviewed literature, and by government entities such as NIOSH and OSHA, have shown that adverse occupational health outcomes are associated with WAG exposures. Recently, a systematic review came to a logical conclusion that evidence for adverse effects of volatile anesthetics on exposed personnel is scarce and inconsistent, but there is no evidence of adverse effects when environmental levels are kept within legal threshold values.<sup>13</sup> It is recognized that further epidemiologic and exposure assessment studies are needed to improve our knowledge and understanding of the effects of occupational exposure to WAGs. New surveillance methods that include systematic data collection, clinical signs, and biomarkers of exposure, may be helpful to formulate consistent and reproducible surveillance criteria for exposed personnel.

Based on a review of the literature, ASPAN presents the following list of potential effects.

- Health effects of exposure to WAGs may cause the following:
  - Headache<sup>1,14,15</sup>
  - Irritability<sup>1,14,15</sup>
  - Fatigue<sup>14,15</sup>
  - Nausea<sup>1,14,15</sup>
  - Drowsiness<sup>1</sup>
  - Depression<sup>1</sup>
  - Difficulties with judgment, audiovisual processing, and coordination<sup>1,16-19</sup>
  - Liver disease<sup>1,16,20</sup>
  - Kidney disease<sup>1,16,20</sup>
  - Malignancies of the lymph or immune system<sup>1,21,22</sup>
  - Impaired B12 metabolic status or pernicious anemia<sup>23-25</sup>
  - Miscarriages<sup>2,18,20,26-30</sup>
  - Congenital abnormalities<sup>2,15,21,27</sup>
  - Low birth weights<sup>7,20,29,30</sup>
  - Exposed spouse's miscarriages and birth defects<sup>2,20,27</sup>
  - Reduced fertility<sup>19,31</sup>
  - Genetic, DNA damage<sup>20,32-35</sup>
  - Cancer<sup>1,11,30,36</sup>
  - Bone marrow depression<sup>17,18,20</sup>
  - Increased homocysteine levels<sup>18</sup>
  - General health and genotoxic risks<sup>37</sup>
  - Changes in antioxidant status<sup>32,34,35,38</sup>

In addition, health effects of surgical anesthetic gases in the elderly contribute to postoperative cognitive decline/dysfunction.<sup>39-42</sup> Health effects of repeated surgical anesthetic exposure in the very young (less than 36 months) are currently being investigated to determine if the exposure may contribute to learning disabilities and autism.<sup>43-54</sup>

## 2. Regulatory Approach

ASPAN and AIHA support the development of engineering controls to decrease exposure to WAGs reduced to the lowest feasible level in the PACU.<sup>7</sup> While the use of prospective studies and carefully designed research protocols is encouraged to elucidate areas of controversy, a responsible approach to worker health and safety dictates that any exposure to waste and trace gases should be kept to the lowest practical level.<sup>2</sup> OSHA states, "the preferred and most effective means of protecting workers is to prevent hazards entering their breathing zone in the first place."<sup>2,3</sup> When workplace exposure limits are implemented by using measures to reduce occupational exposure, few adverse health effects from un-scavenged WAGs have been documented.<sup>8</sup>

Proposed engineering controls to include:

- Market-available source-control scavenging systems which can contain and control WAGs from the exhaled breath of patients in the PACU. In addition, these scavenging systems may help contain and control capture pathogens, such as severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), the pathogen that causes COVID-19, in addition to WAGs<sup>12,17,55-62</sup>
- Controls for the exhaled breath of immediate postoperative patients on ventilators<sup>10,11,55,63</sup>
- Increased heating, ventilation, and air conditioning (HVAC) air exchanges in the PACUs from 6 air changes per hour to 15, similar to the operating rooms<sup>10,11,19,55</sup>
- Regular HVAC monitoring of air quality to be preferably completed during the peak of daily patient activity in the PACU and within the patient breathing zone where the nurses work closely with patients<sup>10,11,55,57,64,68</sup>
- Following OSHA's General Duty Clause<sup>d</sup> for WAGs in PACU until such time as federal standards are established and enacted<sup>2,3,69</sup>
- Adherence to NIOSH Recommended Exposure Limits (RELs) for WAGs<sup>3,11,19</sup>
- Provision of negative airflow rooms for bi-pap and aerosol-generating procedures that induce expired and aerosolized breath/air<sup>3,10,11,59</sup>
- Routine maintenance of these systems<sup>3,11</sup>

### 3. Continued Research

There is a significant and growing body of knowledge on the relationship between the air quality in the PACU and poor health outcomes for healthcare providers in this environment. There may be moderating or predisposing factors that interact with occupational exposure.<sup>11,43,46,59,69</sup> Although exposure to WAGs in the OR has been well documented, there lacks a large body of work for the PACU. This includes the fact that there is little to no surveillance or measurement of air quality (and WAGs source) during peak times of patients' presence in the PACU.<sup>11</sup> Thus, making healthcare providers aware of these risks by utilizing these engineering and administrative controls along with regulatory approaches can lead to a responsible approach for worker health and safety that keeps any exposure to WAGs to the lowest possible practical level.

### Approval of Statement

The statement was developed by ASPAN, supported by the American Industrial Hygiene Association and endorsed by the AIHA Board of Directors vote February 27, 2022. This statement was endorsed by a vote of the ASPAN Board of Directors on April 6, 2022, in Philadelphia, Pennsylvania, and approved by a vote of the ASPAN Representative Assembly on April 7, 2022.

*<sup>d</sup>The General Duty Clause is the portion of the OSHA Act, section 5(a)(1) indicating the requirement of employers to provide a safe workplace environment that is free from hazards.<sup>2,3,69</sup>*

## References

1. Occupational Safety and Health Administration. OSHA Technical Manual (OTM) Section VI: Chapter 1. Hospital Investigations: Health Hazards. Washington, DC: United States Department of Labor; 1991. Accessed May 9, 2022. [https://www.osha.gov/otm/section-6-health-care-facilities/chapter-1#app\\_vi:1](https://www.osha.gov/otm/section-6-health-care-facilities/chapter-1#app_vi:1)
2. Occupational Safety and Health Administration. Anesthetic gases: guidelines for workplace exposures. Washington, DC: United States Department of Labor; 1999. Revised May 18, 2000. Accessed May 9, 2022. <https://www.osha.gov/waste-anesthetic-gases/workplace-exposures-guidelines>
3. National Institute for Occupational Safety and Health. Waste Anesthetic Gases: Occupational Hazards in Hospitals. Publication No. 2007-151. Cincinnati, OH: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention; 2007. Accessed May 9, 2022. <https://www.cdc.gov/niosh/docs/2007-151/pdfs/2007-151.pdf>
4. Kharasch ED. Biotransformation of sevoflurane. *Anesth Analg*. 1995;81(6 Suppl):S27-38. <https://doi.org/10.1097/00000539-199512001-00005>
5. Sherman J, Le C, Lamers V, Eckelman M. Life cycle greenhouse gas emissions of anesthetic drugs. *Anesth Analg*. 2012;114(5):1086-1090. <https://doi.org/10.1213/ANE.0b013e31824f6940>
6. Varughese S, Ahmed R. Environmental and occupational considerations of anesthesia: a narrative review and update. *Anesth Analg*. 2021;133(4):826-835. <https://doi.org/10.1213/ANE.0000000000005504>
7. American Industrial Hygienist Association, American Society of PeriAnesthesia Nurses. Recognition, evaluation and control of waste anesthetic gases in the post-anesthesia care unit. Accessed July 7, 2022. <https://www.aspan.org/Publications-Resources/WAG-White-Paper-and-Presentation>
8. Janssen JE. Ventilation for acceptable indoor air quality. *ASHRAE J*. 1989;31:40-48.
9. Hiller KN, Altamirano AV, Cai C, Tran SF, Williams GW. Evaluation of waste anesthetic gas in the postanesthesia care unit within the patient breathing zone. *Anesthesiol Res Pract*. 2015. <https://doi.org/10.1155/2015/354184>
10. McGlothlin JD, Moenning JE, Cole SS. Evaluation and control of waste anesthetic gases in the postanesthesia care unit. *J Perianesth Nurs*. 2014;29(4):298-312. <https://doi.org/10.1016/j.jopan.2013.09.010>
11. Moenning JE, Krenzischek DA, McGlothlin JD. Chapter 13: Waste anesthetic gases (WAGs): minimizing health risks and increasing awareness. In: Gross KA, ed. *Advanced Practice and Leadership in Radiology Nursing*. Springer; 2020:149-159.
12. Tallent R, Corcoran J, Sebastian J. Evaluation of a novel waste anaesthetic gas scavenger device for use during recovery from anaesthesia. *Anaesthesia*. 2018;73(1):59-64. <https://doi.org/10.1111/anae.14100>
13. Molina Aragonés JM, Ayora AA, Ribalta AB, et al. Occupational exposure to volatile anaesthetics: a systematic review. *Occup Med*. 2016;66(3):202-207. <https://doi.org/10.1093/occmed/kqv193>
14. Rogers B. Exposure to waste anesthetic gases--a review of toxic effects. *AAOHN J*. 1986;34(12):574-579.
15. Vaisman AI. Working conditions in the operating room and their effect on the health of anesthesiologists. *Eksp Khir Anestheziol*. 1967;12(3):44-49.
16. Bruce DL, Bach MJ, Arbit J. Trace anesthetic effects on perceptual, cognitive, and motor skills. *Anesthesiology*. 1974;40:453-457.
17. National Institute for Occupational Safety and Health (NIOSH). Control of nitrous oxide in dental operatories. Cincinnati, OH: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention; 2007. Accessed May 9, 2022. <https://www.cdc.gov/niosh/docs/hazardcontrol/hc3.html>
18. Sanders RD, Weimann J, Maze M, Warner DS, Warner MA. Biologic effects of nitrous oxide: a mechanistic and toxicologic review. *Anesthesiology*. 2008;109(4):707-722. <https://doi.org/10.1097/ALN.0b013e3181870a17>
19. Sessler DI, Badgwell JM. Exposure of postoperative nurses to exhaled anesthetic gases. *Anesth Analg*. 1998;87(5):1083-1088. <https://doi.org/10.1097/00000539-199811000-00019>
20. Cohen EN, Gift HC, Brown BW, et al. Occupational disease in dentistry and chronic exposure to trace anesthetic gases. *J Am Dent Assoc*. 1980;101(1):21-31. <https://doi.org/10.14219/jada.archive.1980.0345>
21. Bruce DL, Eide KA, Linde HW, Eckenhoff JE. Causes of death among anesthesiologists: a 20-year survey. *Anesthesiology*. 1968;29(3):565-569. <https://doi.org/10.1097/00000542-196805000-00039>

22. Bruce DL, Eide KA, Smith NJ, Seltzer F, Dykes MH. A prospective survey of anesthesiologist mortality, 1967–1971. *Anesthesiology*. 1974;41(1):71-74. <https://doi.org/10.1097/0000542-197407000-00017>
23. Amess JA, Burman JF, Rees GM, Nancekieveill DG, Mollin DL. Megaloblastic haemopoiesis in patients receiving nitrous oxide. *Lancet*. 1978;12(2):339-342. [https://doi.org/10.1016/s0140-6736\(78\)92941-0](https://doi.org/10.1016/s0140-6736(78)92941-0)
24. Banks RG, Henderson RJ, Pratt JM. Reactions of gases in solution. Part III. Some reactions of nitrous oxide with transition-metal complexes. *J Chem Soc A*. 1968;2886-2889. <https://doi.org/10.1039/J19680002886>
25. Krajewski W, Kucharska M, Pilacik B, et al. Impaired vitamin B12 metabolic status in healthcare workers occupationally exposed to nitrous oxide. *Br J Anaesth*. 2007;99(6):812–8. <https://doi.org/10.1093/bja/aem280>
26. Buring JE, Hennekens CH, Mayrent SL, Rosner B, Greenberg ER, Colton T. Health experiences of operating room personnel. *Anesthesiology*. 1985;62(3):325-330. <https://doi.org/10.1097/0000542-198503000-00018>
27. Cohen EN, Bellville JW, Brown Jr BW. Anesthesia, pregnancy, and miscarriage: a study of operating room nurses and anesthesiologists. *Anesthesiology*. 1971;35(4):343-347. <https://doi.org/10.1097/0000542-197110000-00005>
28. Corbett TH, Cornell RG, Endres JL, Lieding K. Birth defects among children of nurse-anesthetists. *Anesthesiology*. 1974;41(4):341-344.
29. Pharoah POD, Alberman E, Doyle P, Chamberlain G. Outcome of pregnancy among women in anaesthetic practice. *Lancet*. 1977;309(8001):34-36. [https://doi.org/10.1016/s0140-6736\(77\)91666-x](https://doi.org/10.1016/s0140-6736(77)91666-x)
30. Tomlin PJ. Health problems of anaesthetists and their families in the West Midlands. *Br Med J*. 1979;1(6166):779-784. <https://doi.org/10.1136/bmj.1.6166.779>
31. Rowland AS, Baird DD, Weinberg CR, Shore DL, Shy CM, Wilcox AJ. Reduced fertility among women employed as dental assistants exposed to high levels of nitrous oxide. *N Engl J Med*. 1992;327(14):993-997. <https://doi.org/10.1056/NEJM199210013271405>
32. Casale T, Caciari T, Rosati MV, et al. Anesthetic gases and occupationally exposed workers. *Environ Toxicol Pharmacol*. 2014;37(1):267-274. <https://doi.org/10.1016/j.etap.2013.12.003>
33. Lucio LMC, Braz MG, Nascimento Jr PD, Braz JRC, Braz LG. Occupational hazards, DNA damage, and oxidative stress on exposure to waste anesthetic gases. *Braz J Anesthesiol*. 2018;68(1):33-41. <https://doi.org/10.1016/j.bjan.2017.05.001>
34. Costa Paes ER, Braz MG, Lima JT, et al. DNA damage and antioxidant status in medical residents occupationally exposed to waste anesthetic gases. *Acta Cir Bras*. 2014;29(4):280-286. <https://doi.org/10.1590/s0102-86502014000400010>
35. Rocha TL, Dias-Junior CA, Possomato-Vieira JS, et al. Sevoflurane induces DNA damage whereas isoflurane leads to higher antioxidative status in anesthetized rats. *Biomed Res Int*. 2015;2015:264971. <https://doi.org/10.1155/2015/264971>
36. Corbett TH. Cancer and congenital anomalies associated with anesthetics. *Ann N Y Acad Sci*. 1976;271:58-66. <https://doi.org/10.1111/j.1749-6632.1976.tb23093.x>
37. Fodale V, Mondello S, Aloisi C, Schifilliti D, Santamaria LB. Genotoxic effects of anesthetic agents. *Expert Opin Drug Saf*. 2008;7(4):447-458. <https://doi.org/10.1517/14740338.7.4.447>
38. Deng HB, Li FX, Cai YH, Xu SY. Waste anesthetic gas exposure and strategies for solution. *J Anesth*. 2018;32(2):269-282. <https://doi.org/10.1007/s00540-018-2448-1>
39. Bittner EA, Yue Y, Xie Z. Brief review: anesthetic neurotoxicity in the elderly, cognitive dysfunction and Alzheimer’s disease. *Can J Anaesth*. 2011;58(2):216-223. <https://dx.doi.org/10.1007%2Fs12630-010-9418-x>
40. Craddock TJA, Kurian P, Preto J, et al. Anesthetic alterations of collective terahertz oscillations in tubulin correlate with clinical potency: implications for anesthetic action and post-operative cognitive dysfunction. *Sci Rep*. 2017;7(1):1-12. <https://doi.org/10.1038/s41598-017-09992-7>
41. Rundshagen I. Postoperative cognitive dysfunction. *Dtsch Arztebl Int*. 2014;111(8):119-125. <https://doi.org/10.3238/arztebl.2014.0119>
42. Wang W, Wan Y, Wu H, et al. Postoperative cognitive dysfunction: current developments in mechanism and prevention. *Med Sci Monit*. 2014;20:1908-1912. <https://doi.org/10.12659/MSM.892485>
43. Card EB, Wells NL. An introduction to the Smart Tots consensus statement on the use of anesthetic and sedative drugs in infants and toddlers. *J Perianesth Nurs*. 2016;31(1):3-10. <https://doi.org/10.1016/j.jopan.2015.12.001>

44. Chinn GA, Russell JMS, Sall JW. Is a short anesthetic exposure in children safe? Time will tell: a focused commentary of the GAS and PANDA trials. *Ann Transl Med.* 2016;4(20):408. <https://doi.org/10.21037/atm.2016.10.43>
45. Feng YP, Yang TS, Chung CH, Chien WC, Wong CS. Early childhood general anesthesia exposure associated with later developmental delay: a national population-based cohort study. *PLoS One.* 2020;15(9):e0238289. <https://doi.org/10.1371/journal.pone.0238289>
46. Flick RP, Katusic SK, Colligan RC, et al. Cognitive and behavioral outcomes after early exposure to anesthesia and surgery. *Pediatrics.* 2011;128(5), e1053-e1061. <https://doi.org/10.1542/peds.2011-0351>
47. Flick RP, Ing CH. SmartTots and FDA: anesthetic-related neurotoxicity in children—ASA works in close cooperation with FDA and others to advance research. *ASA Newsletter.* 2015;79(3):14-16.
48. Gleich SJ, Shi Y, Flick R, et al. Hypotension and adverse neurodevelopmental outcomes among children with multiple exposures to general anesthesia: subanalysis of the Mayo Anesthesia Safety in Kids (MASK) Study. *Paediatr Anaesth.* 2021;31(3):282-289. <https://doi.org/10.1111/pan.14106>
49. Ing CH, Flick RP. Anesthetic-related neurotoxicity—current state of the art. *ASA Monitor.* 2016;80(11):16-18.
50. O’Leary JD, Janus M, Duku E, et al. A population-based study evaluating the association between surgery in early life and child development at primary school entry. *Anesthesiology.* 2016;125(2):272-279. <https://doi.org/10.1097/ALN.0000000000001200>
51. Raper J, Kamat PP. Pediatric Anesthetic and Sedation Neurotoxicity in the Developing Brain. In: Kamat PP, Berkenbosch JW, eds. *Sedation and Analgesia for the Pediatric Intensivist.* Springer; 2021:233-244. [https://doi.org/10.1007/978-3-030-52555-2\\_17](https://doi.org/10.1007/978-3-030-52555-2_17)
52. Orser BA, Evers AS, Suresh S, Hertz S. The GAS trial. *Lancet.* 2016;387(10028):1613-1614. [https://doi.org/10.1016/S0140-6736\(16\)30199-4](https://doi.org/10.1016/S0140-6736(16)30199-4)
53. Tanzi MG. New warning added to labels of general anesthetic and sedation agents. *Pharmacy Today.* 2017;23(7):2.
54. Vutskits L, Culley DJ. GAS, PANDA, and MASK: no evidence of clinical anesthetic neurotoxicity! *Anesthesiology.* 2019;131(4):762-764. <https://doi.org/10.1097/ALN.0000000000002863>
55. American Society of Perianesthesia Nurses. A position statement on air safety in the perianesthesia environment. *J Perianesth Nurs.* 1996;11(4):203–5.
56. Applegate B. A laboratory study of a patient scavenging system (part II). *Conference Session of the American Society of PeriAnesthesia Nurses.* United States. April 25-29, 2021.
57. Badgwell JM. A clinical evaluation of an operational postanesthesia care unit source control system. *J Perianesth Nurs.* 1997;12(2):73-81.
58. Centers for Disease Control and Prevention. Infection prevention and control recommendations for hospitalized patients under investigation for Ebola virus disease in U.S. hospitals. 2010. <https://www.cdc.gov/vhf/ebola/clinicians/evd/infection-control.html>
59. Horton JL. Laboratory study of a scavenging mask system to evaluate and control airborne pathogens for healthcare workers in the Post Anesthesia Care Unit (PACU) and Intensive Care Unit (ICU). *Open Access Theses.* 2014;1027. [https://docs.lib.purdue.edu/cgi/viewcontent.cgi?article=1763&context=open\\_access\\_theses&httpsredir=1](https://docs.lib.purdue.edu/cgi/viewcontent.cgi?article=1763&context=open_access_theses&httpsredir=1)
60. McGlothlin JD. A laboratory study of a patient scavenging system (part I). *Conference Session of the American Society of PeriAnesthesia Nurses.* United States. April 25-29, 2021.
61. Rademaker AM, McGlothlin JD, Moenning JE, Bagnoli M, Carlson G, Griffin C. Evaluation of two nitrous oxide scavenging systems using infrared thermography to visualize and control emissions. *J Am Dent Assoc.* 2009;140(2):190-199. <https://doi.org/10.14219/jada.archive.2009.0133>
62. Sedgley C, Applegate B, Nagel A, Hall D. Real-time imaging and quantification of bioluminescent bacteria in root canals in vitro. *J Endod.* 2004;30(12):893-898. <https://doi.org/10.1097/01.don.0000132299.02265.6c>
63. Brodsky JB. Exposure to anesthetic gases: a controversy. *AORN J.* 1983;38(1):132-144. [https://doi.org/10.1016/s0001-2092\(07\)69556-8](https://doi.org/10.1016/s0001-2092(07)69556-8)
64. Austin PR, Austin PJ. Measurement of nitrous oxide concentrations in a simulated post anesthesia care unit environment. *J Perianesth Nurs.* 1996;11(4):259-266. [https://doi.org/10.1016/s1089-9472\(96\)80026-8](https://doi.org/10.1016/s1089-9472(96)80026-8)



65. Cheung SK, Özelsel T, Rashiq S, Tsui BC. Postoperative environmental anesthetic vapour concentrations following removal of the airway device in the operating room versus the postanesthesia care unit. *Can J Anaesth*. 2016;63(9):1016-1021. <https://doi.org/10.1007/s12630-016-0678-y>
66. Cope KA, Merritt WT, Krenzischek DA, et al. Phase II collaborative pilot study: preliminary analysis of central neural effects from exposure to volatile anesthetics in the PACU. *J Perianesth Nurs*. 2002;17(4):240-250. <https://doi.org/10.1053/jpan.2002.34167>
67. Krenzischek DA, Schaefer J, Nolan M, et al. Phase I collaborative pilot study: waste anesthetic gas levels in the PACU. *J Perianesth Nurs*. 2002;17(4):227-239. <https://doi.org/10.1053/jpan.2002.34166>
68. Summer G, Lirk P, Hoerauf K, et al. Sevoflurane in exhaled air of operating room personnel. *Anesth Analg*. 2003;97(4):1070-1073. <https://doi.org/10.1213/01.ANE.0000081796.67539.27>
69. National Institute for Occupational Safety and Health. OSHA's General Duty Clause. Reviewed February 7, 2020. Accessed July 15, 2022. [https://www.cdc.gov/WPVHC/Nurses/Course/Slide/Unit5\\_4](https://www.cdc.gov/WPVHC/Nurses/Course/Slide/Unit5_4)

## Additional Reading

Aggarwal S, Axelrod D, Bell C, et al. Greening the operating room and perioperative arena: environmental sustainability for anesthesia practice. 2015. Accessed June 2, 2020. <https://www.asahq.org/about-asa/governance-and-committees/asa-committees/environmental-sustainability/greening-the-operating-room-complete>

American Society of Anesthesiologists. Task Force on Trace Anesthetic Gases. Waste anesthetic gases: information for management in anesthetizing areas and the postanesthesia care unit (PACU). ASA; 1999. [https://www.cdc.gov/niosh/docket/archive/pdfs/niosh-064/0064-010199-mcgregor\\_ref2.pdf](https://www.cdc.gov/niosh/docket/archive/pdfs/niosh-064/0064-010199-mcgregor_ref2.pdf)

Hönemann C, Hagemann O, Doll D. Inhalational anaesthesia with low fresh gas flow. *Indian J Anaesth*. 2013;57(4):345-350. <https://dx.doi.org/10.4103%2F0019-5049.118569>

Krapez JR, Saloojee Y, Hinds CJ, Hackett GH, Cole PV. Blood concentrations of nitrous oxide in theatre personnel. *Br J Anaesth*. 1980;52(11):1143-1148. <https://doi.org/10.1093/bja/52.11.1143>

National Institute for Occupational Safety and Health. Respiratory protective devices. Code of Federal Regulations. 1995. Title 42, Part 84.

Peter MA, Malherbe S. Environmentally sustainable perioperative medicine: simple strategies for anesthetic practice. *Can J Anaesth*. 2020;67(8):1044-1063. <https://doi.org/10.1007/s12630-020-01726-0>

World Resources Institute. World Greenhouse Gas Emissions. 2016. Accessed <https://www.wri.org/resources/data-visualizations/world-greenhouse-gas-emissions-2016>